

BINARY ADDITIVE PROBLEMS: THEOREMS OF LANDAU AND HARDY-LITTLEWOOD TYPE

VLADIMIR SHEVELEV

ABSTRACT. We prove theorems of Landau and Hardy-Littlewood type for Goldbach, Chen, Lemoine-Levy and other binary partitions of positive integers. We also pose some new conjectures.

1. INTRODUCTION

Let P be the set of primes. For an even number $n \geq 6$, let $n = p + q$ with $p, q \in P$ be a Goldbach partition of n . Denote $g(n)$ the number of the unordered Goldbach partitions of n . Denote, furthermore, as in [11], $N_2(n)$ the number of such partitions with the taking into account the order of parts. Then, evidently,

$$(1) \quad N_2(n) = \begin{cases} 2g(n), & \text{if } n/2 \notin P, \\ 2g(n) - 1, & \text{if } n/2 \in P. \end{cases}$$

Justifying the posing of their famous conjecture

$$(2) \quad N_2(n) \sim 2C_{tw}n / \ln^2 n \prod_{p|n, \ p \text{ odd prime}} \frac{p-1}{p-2},$$

where C_{tw} is the "twin constant"

$$C_{tw} = \prod_{p \text{ odd prime}} \frac{p(p-2)}{(p-1)^2} = 0,66016\dots,$$

Hardy and Littlewood [11, pp.35-36] reproved the following Landau asymptotics

$$(3) \quad \sum_{i=1}^n N_2(2i) \sim n^2 / (2 \ln^2 n).$$

In fact, using (3), they obtained the following result.

Theorem 1. *If limit*

$$\lim_{\text{even } x \rightarrow \infty} \left(\frac{N_2(x) \ln^2 x}{x} \prod_{p|x, \ p \text{ odd prime}} \frac{p-2}{p-1} \right)$$

exists, then it equals to $2C_{tw}$.

Landau proved (3) using his identity for the φ -function which he obtained in [13]:

$$(4) \quad \sum_{d|n} \mu^2(d)/\varphi(d) = n/\varphi(n),$$

while Hardy and Littlewood proved (3) using their tauberian theorem [10]. A modern interesting information about Landau formulas (3),(4) one can find in [8],[12]. In 1998, Dusart [5] obtained the following excellent estimates for the prime counting function $\pi(x)$: if $x \geq 355991$, then

$$(5) \quad x/\ln x + x/\ln^2 x < \pi(x) < x/\ln x + x/\ln^2 x + 2.51x/\ln^3 x.$$

He also obtained the best lower estimate for n th prime p_n [6], such that the modern known estimates for p_n are ([1]):

$$(6) \quad n \ln n + n \ln \ln n - n < p_n < n \ln n + n \ln \ln n, \quad n \geq 6.$$

These estimates are much stronger than those ones which could be obtained from the well known estimates of Rosser [16]. In this paper, using our identities [17] and estimates (5)-(6), we give a new proof of (3) and close results for sums of numbers of Lemoine-Levy, Chen and some other known binary partitions; we also give new theorems like Theorem 1 (Theorems 4 and 6) and propose a new short way for proof of theorems of such type.

2. GOLDBACH PARTITIONS

Let P_1 be the set of odd primes and $\pi_1(x)$ be its counting function. In [17, example 1] we showed that, for $x = 2n$,

$$(7) \quad \sum_{i=1}^{x/2} g(2i) = \sum_{3 \leq p \leq x/2, p \in P} \pi_1(x-p) - \binom{\pi_1(x/2)}{2}.$$

Using (5), we find

$$\begin{aligned} \sum_{i=1}^{x/2} g(2i) &= \frac{1}{\ln x} \sum_{3 \leq p \leq x/2, p \in P} (x-p) - \frac{x^2}{8 \ln^2 x} + O\left(\frac{x^2}{\ln^3 x}\right) \\ (8) \quad &= \frac{x}{\ln x} \pi_1(x/2) - \frac{x^2}{8 \ln^2 x} - \frac{1}{\ln x} \sum_{i=2}^{\pi(x/2)} p_i + O\left(\frac{x^2}{\ln^3 x}\right) \\ &= \frac{3x^2}{8 \ln^2 x} - \frac{1}{\ln x} \sum_{i=2}^{\pi(x/2)} p_i + O\left(\frac{x^2}{\ln^3 x}\right). \end{aligned}$$

Furthermore, by (5)-(6), we have

$$(9) \quad \sum_{i=2}^{\pi(x/2)} p_i = \sum_{i=2}^{\pi(x/2)} i \ln i + (\ln \ln x) \pi^2(x/2)/2 + O\left(\frac{x^2}{\ln^2 x}\right).$$

Finally, notice that

$$(10) \quad \sum_{i=2}^{\pi(x/2)} i \ln i = \int_2^{\pi(x/2)} t \ln t dt + O(\pi(x/2) \ln \pi(x/2)) = \frac{x^2}{8 \ln x} + O\left(\frac{x^2}{\ln^2 x}\right).$$

Now from (8)-(10) we obtain theorem Landau in the form:

Theorem 2.

$$(11) \quad \sum_{i=1}^{x/2} g(2i) = \frac{x^2}{4 \ln^2 x} + O\left(\frac{x^2 \ln \ln x}{\ln^3 x}\right).$$

Note that (3) follows from (11) and (1). ■

Corollary 1.

$$\sum_{p+q \leq x, \substack{p \leq q, \\ p, q \in P}} \ln(p+q) = \frac{x^2}{4 \ln x} + O\left(\frac{x^2 \ln \ln x}{\ln^2 x}\right).$$

Proof. Putting $x = 2n$, we prove that

$$\sum_{p+q \leq 2n, \substack{p \leq q, \\ p, q \in P}} \ln(p+q) = \frac{n^2}{\ln n} + O\left(\frac{n^2 \ln \ln n}{\ln^2 n}\right).$$

Denoting, for $n \geq 3$,

$$G(n) = \sum_{i=3}^n g(2i)$$

and using the summation by parts, we have

$$\sum_{k=3}^n g(2k+2) \ln(2k) = G(n+1) \ln(2n+2) - G(3) \ln 6 - \sum_{k=3}^n G(n+1) \ln\left(1 + \frac{1}{k}\right).$$

Because of, by Theorem 2, $G(n)/n$ is increasing since $n \geq n_0$, then we have

$$\begin{aligned} \sum_{k=3}^n G(n+1) \ln\left(1 + \frac{1}{k}\right) &\leq \sum_{k=3}^n G(k+1)/k \leq \\ &\sum_{k=3}^n G(n+1)/n + O(1) = G(n+1) + O(1). \end{aligned}$$

Thus, by Theorem 2,

$$\sum_{k=3}^n g(2k+2) \ln(2k) = n^2 / \ln n + O\left(\frac{n^2 \ln \ln n}{\ln^2 n}\right)$$

and sufficiently to notice that the following identity is valid

$$\prod_{p+q \leq x, \ p \leq q, \ p, \ q \in P} (p+q) = \prod_{k=3}^n (2k)^{g(2k)}. \blacksquare$$

3. GRANVILLE-LUNE-REELE PARTITIONS

In 1989, A. Granville, H. te Riele and J. van de Lune [9] conjectured that if, for even n , $p = p(n)$ means the least prime such that $n - p$ is also prime, then

$$(12) \quad p(n) = O(\ln^2 n \ln \ln n)$$

In connection with this, a Golgbach partition of n , for the least part of which (12) is valid (for the minimal constant C in $O(\dots)$), we call a Granville-Lune-Reele partition (GLR-partition) of n . Denote $\nu(n)$ the number of unordered GLR-partition of n . According to (7), we have

$$(13) \quad \sum_{i=1}^{x/2} \nu(2i) = \sum_{3 \leq p \leq C \ln^2 x \ln \ln x, \ p \in P} \pi_1(x - p) + O(1),$$

where $O(1)$ corresponds to the finite set of even n having a GLR-partition of the form $n = n/2 + n/2$. As in above, now we find

$$(14) \quad \sum_{i=1}^{x/2} \nu(2i) = \frac{C}{2} x \ln x + O(x).$$

By GLR-conjecture (12), for every sufficiently large n , $\nu(n) > 0$. On the other hand, $\nu(n) \leq C \ln^2 n \ln \ln n$. The following arguments show that the GLR-conjecture is essentially unprovable. Note that the number of GLR-partitions of 2^k , $k \geq 3$ is not more than $C_1 k^2 \ln k$, such that the total number of such partitions for $2^k \leq x$ is $O(\ln^3 x \ln \ln x)$. Note, furthermore, that, the most primes of Goldbach partitions of 2^k are in interval $(2^{k-1}, 2^k)$ and do not take part in Goldbach partitions as the most parts of neither previous nor following powers (see sequence A152451 in [18]). Thus, if to remove the most primes in every GLR-representations of $2^k, k = 3, 4, \dots$, we obtain a sequence P^* of primes with the counting function $\pi^*(x) = \pi(x) - O(\ln^3 x \ln \ln x)$. Replacing in P^* the first N terms by N consecutive primes (with arbitrary large N), we obtain a sequence which essentially is indistinguishable from the sequence P of all primes with help of the approximation of $\pi(x)$ by $li(x)$, since, according to well known Littlewood result [14], the remainder term in the theorem of primes could not be less than $\sqrt{x} \log \log \log x / \log x$. But, according to our construction, for this sequence of primes we have infinitely many even numbers for which the

GLR-conjecture is wrong. Thus the conjecture is essentially unprovable.

Remark 1. Note that if $A(x)$ is the counting function of sequence A152451 [18] then, because of $\sum_{3 \leq n \leq \log x} 2^n/n^2 = O(x/\ln^2 x)$, we have $A(x) = \pi(x) - O(x/\ln^2 x)$. It is well known that for the approximation of $\pi(x)$ by $x/\ln x$ the remainder term is, as the best, $O(x/\ln^2 x)$. Therefore, beginning the constructing algorithm of sequence A152451 from $k \geq N$ (with arbitrary large N) we obtain a sequence which essentially is indistinguishable from the sequence of all odd primes with help of the approximation of $\pi(x)$ by $x/\ln x$, although, according to the construction, we have infinitely many even numbers which are not expressible by sum of two primes of this sequence. Hence, in principle, it is impossible to prove the binary Goldbach conjecture using such approximation of $\pi(x)$.

Notice also that, a more economical algorithm leads to sequence A156284 [18] (see also an interest sequence A156537 of "wells").

4. LEMOINE-LEVY PARTITIONS

Let $l(n)$, odd $n \geq 1$, denote the number of decompositions of n into unordered sums of a prime and a doubled prime (Lemoine-Levy partitions). Then, putting in [17, example 3] $2n - 1 = x$, we have:

$$\begin{aligned} \sum_{i=1}^{\frac{x+1}{2}} l(2i-1) &= \sum_{2 \leq p \leq \frac{x+1}{2}, p \in P} \pi\left(\frac{x-p}{2}\right) + \sum_{2 \leq p \leq \frac{x+1}{4}, p \in P} \pi(x-2p) \\ (15) \quad &\quad - \pi\left(\frac{x+1}{2}\right) \pi\left(\frac{x+1}{4}\right). \end{aligned}$$

Now, as in Section 2, using (5)-(6), we obtained the result which shows that in average $g(n)$ and $l(n)$ asymptotically coincide.

Theorem 3.

$$(16) \quad \sum_{i=1}^{\frac{x+1}{2}} l(2i-1) = \frac{x^2}{4 \ln^2 x} + O\left(\frac{x^2 \ln \ln x}{\ln^3 x}\right).$$

Corollary 2.

$$\sum_{p+2q \leq x, p, q \in P} \ln(p+2q) = \frac{x^2}{4 \ln x} + O\left(\frac{x^2 \ln \ln x}{\ln^2 x}\right).$$

Proof repeats proof of Corollary 1 if to notice that

$$\prod_{p+2q \leq 2n-1, p, q \in P} (p+2q) = \prod_{k=5}^n (2k-1)^{l(2k-1)}. \blacksquare$$

Moreover, using the scheme of proof of Theorem 6 (see below), it could be proved the following result which is similar to Theorem 1.

Theorem 4. *If limit*

$$\lim_{\substack{\text{odd} \\ x \rightarrow \infty}} \left(\frac{l(x) \ln^2 x}{x} \prod_{p|x, \ p \text{ odd prime}} \frac{p-2}{p-1} \right)$$

exists, then it equals to C_{tw} .

This theorem is a base for a very close to (2) conjecture: for odd $n \rightarrow \infty$

$$(17) \quad l(n) \sim C_{tw} n / \ln^2 n \prod_{p|n, \ p \text{ odd prime}} \frac{p-1}{p-2}.$$

Remark 2. *If to consider the Lemoine-Levy's partitions with small least parts, as in Section 3, then one can use the similar constructive arguments (see comments to our sequences A152460 – A152461 in [18]).*

5. CHEN PARTITIONS

Let $c(n)$ denote the number of decompositions of even n into prime and prime or semiprime ([19]). In 1966 Chen [2] did a very important step towards Goldbach conjecture, proving that for sufficiently large n

$$(18) \quad c(n) \geq a C_{tw} \left(\prod_{p|n, \ p \text{ odd prime}} \frac{p-1}{p-2} \right) n / \ln^2 n,$$

where $a = 0.098$ [2] with improvements $a = 0.67$ [3] and 1.0974 [4]. Ross [15] gave a large simplification of Chen's proof.

Putting in [17, example 2] $2n = x$ and $g_1(2i) + g_2(2i) = c(2i)$, $i = 1, \dots, x/2$, we have:

$$(19) \quad \sum_{i=1}^{x/2} c(2i) = \sum_{9 \leq q \leq x/2, \ q \in P_{2,1}} \pi_1(x-q) + \sum_{3 \leq p \leq x/2, \ p \in P} \pi_{2,1}(x-p) \\ - \pi_1(x/2)(\pi_{2,1}(x/2) + 1/2(\pi_1(x/2) - 1)) + \pi(x/2 - 1) + 1, \ x \geq 4.$$

where $\pi_1(x)$ is the counting function of odd primes and $\pi_{2,1}(x)$ is the counting function of the set $P_{2,1} = \{q_n\}_{n \geq 1}$ of odd semiprimes, such that (sf [19]):

$$(20) \quad \pi_{2,1}(x) = \sum_{3 \leq p \leq \sqrt{x}, \ p \in P} (\pi(x/p) - \pi(p) + 1).$$

Using (5)-(6), we find

$$(21) \quad \sum_{3 \leq p \leq \sqrt{x}, \ p \in P} \pi(x/p) = x \ln \ln x / \ln x + O(x \ln \ln x / \ln^2 x),$$

$$(22) \quad \sum_{3 \leq p \leq \sqrt{x}, p \in P} \pi(p) = O(x / \ln^2 x),$$

such that, by (20), we have

$$(23) \quad \pi_{2,1}(x) = x \ln \ln x / \ln x + O(x \ln \ln x / \ln^2 x).$$

Note that, by (23),

$$n = \pi_{2,1}(q_n) = q_n \ln \ln q_n / \ln q_n + O(q_n \ln \ln q_n / \ln^2 q_n),$$

or

$$(24) \quad q_n = n \ln q_n / \ln \ln q_n + O(n / \ln \ln q_n).$$

Iterating of (24), we find

$$(25) \quad \begin{aligned} q_n &= n \ln q_n / \ln \ln q_n + O(n / \ln \ln q_n) \\ &= \frac{n(\ln n + \ln \ln q_n - \ln \ln \ln q_n + o(1))}{\ln(\ln n + \ln \ln q_n - \ln \ln \ln q_n + o(1))} (1 + o(1)) \end{aligned}$$

$$(26) \quad \ln q_n = \ln n + \ln \ln n (1 + o(1)).$$

Using (25)-(26), we obtain that

$$(27) \quad q_n = n \ln n / \ln \ln n + O(n \ln \ln n).$$

Using (23),(27), from (19) we have

$$(28) \quad \begin{aligned} \sum_{i=1}^{x/2} c(2i) &= \frac{1}{\ln x} \sum_{9 \leq q \leq x/2, q \in P_{2,1}} (x - q) + \frac{\ln \ln x}{\ln x} \sum_{3 \leq p \leq x/2, p \in P} (x - p) \\ &\quad - \frac{x^2 \ln \ln x}{4 \ln^2 x} + O\left(\frac{x^2}{\ln^2 x}\right) \\ &= \frac{3}{4} \frac{x^2 \ln \ln x}{\ln^2 x} - \frac{1}{\ln x} \sum_{i=1}^{\pi_{2,1}(x/2)} q_i - \frac{\ln \ln x}{\ln x} \sum_{i=1}^{\pi(x/2)} p_i + O\left(\frac{x^2}{\ln^2 x}\right), \end{aligned}$$

or, taking into account (9)-(10),

$$(29) \quad \sum_{i=1}^{x/2} c(2i) = \frac{5}{8} \frac{x^2 \ln \ln x}{\ln^2 x} - \frac{1}{\ln x} \sum_{i=1}^{\pi_{2,1}(x/2)} q_i + O\left(\frac{x^2}{\ln^2 x}\right).$$

Furthermore, using (27) and (23), we have

$$\begin{aligned} \sum_{i=1}^{\pi_{2,1}(x/2)} q_i &= \sum_{i=1}^{\pi_{2,1}(x/2)} i \ln i / \ln \ln i + O\left(\sum_{i=1}^{\pi_{2,1}(x/2)} i \ln \ln i\right) \\ &= \int_3^{\pi_{2,1}(x/2)} \frac{t \ln t}{\ln \ln t} dt + O(\pi_{2,1}^2(x/2) \ln \ln \pi_{2,1}(x/2)) \end{aligned}$$

$$(30) \quad = \int_3^{\pi_{2,1}(x/2)} \frac{t \ln t}{\ln \ln t} dt + O\left(\frac{x^2 (\ln \ln x)^3}{\ln^2 x}\right).$$

Finally,

$$(31) \quad \int_3^{\pi_{2,1}(x/2)} \frac{t \ln t}{\ln \ln t} dt = \left(\frac{1}{2}\pi_{2,1}^2(x/2) \ln \pi_{2,1}(x/2) - \frac{1}{4}\pi_{2,1}^2(x/2)\right) \frac{1}{\ln \ln \pi_{2,1}(x/2)} \\ + O\left(\int_3^{\pi_{2,1}(x/2)} \frac{t dt}{(\ln \ln t)^2}\right) = \frac{x^2 \ln \ln x}{8 \ln x} + O\left(\left(\frac{x \ln \ln x}{\ln x}\right)^2\right).$$

Now from (29)-(31) we obtain the following result.

Theorem 5.

$$(32) \quad \sum_{i=1}^{\frac{x}{2}} c(2i) = \frac{x^2 \ln \ln x}{2 \ln^2 x} + O\left(\frac{x^2}{\ln^2 x}\right).$$

Similar to Corollaries 1,2, in view of the identity

$$\prod_{p+q \leq 2n, \ p \in P, \ q \in P_{2,1} \cup P} (p+q) = \prod_{k=3}^n (2k)^{c(2k)}, \blacksquare$$

we obtain the following formula.

Corollary 3.

$$\sum_{p+q \leq x, \ p \in P, \ q \in P_{2,1} \cup P} \ln(p+q) = \frac{x^2 \ln \ln x}{2 \ln x} + O\left(\frac{x^2 \ln \ln x}{\ln^2 x}\right).$$

Furthermore, we show that the following theorem is valid.

Theorem 6. *If limit*

$$\lim_{\text{even } x \rightarrow \infty} \left(\frac{c(x) \ln^2 x}{x \ln \ln x} \prod_{p|x, \ p \text{ odd prime}} \frac{p-2}{p-1} \right)$$

exists, then it equals to $2C_{tw}$.

This theorem is a base for a similar to (2) conjecture: for even $n \rightarrow \infty$

$$(33) \quad c(n) \sim 2C_{tw} n \ln \ln n / \ln^2 n \prod_{p|n, \ p \text{ odd prime}} \frac{p-1}{p-2}.$$

Note that inequality (18) is weaker than (33). Therefore, it is nature to think that a proof of the binary Goldbach conjecture will come about a form of a weaker inequality than (2), for instance,

$$c(n) \geq aC_{tw} \left(\prod_{p|n, \ p \text{ odd prime}} \frac{p-1}{p-2} \right) n / \ln^3 n.$$

6. PROOF OF THEOREM 6

By the following way of proof, one can also reprove Theorem 1 and prove Theorem 4. We use a known statement belonging to O. Stolz (sf [7, pp. 67-68]).

Lemma 1. *Let $y_n \rightarrow \infty$ and be increasing at least for $n \geq n_0$:*

$$\Delta y_n = y_n - y_{n-1} > 0, \quad n \geq n_0.$$

If, for some another variable x_n , there exists limit

$$\lim_{n \rightarrow \infty} \frac{\Delta x_n}{\Delta y_n} = a, \quad |a| \leq \infty,$$

then $\lim_{n \rightarrow \infty} \frac{x_n}{y_n}$ exists as well and equals to a .

Lemma 2. *If, for $a_n > 0$, we have $1/a_n = O(1)$ ($n \rightarrow \infty$) and, for $\varepsilon > 0$,*

$$(34) \quad \sum_{i=1}^n a_i = O(n(\ln n)^{1-\varepsilon}),$$

then

$$(35) \quad \sum_{i=3}^n \frac{\ln \ln i}{\ln^2 i} i a_i \sim \frac{\ln \ln n}{\ln^2 n} \sum_{i=3}^n i a_i.$$

Proof. Putting

$$(36) \quad \lambda(n) = \frac{\ln^2 n}{\ln \ln n}, \quad y_n = \sum_{i=3}^n i a_i, \quad x_n = \lambda(n) \sum_{i=3}^n \frac{1}{\lambda(i)} i a_i,$$

we should prove that $\lim_{n \rightarrow \infty} \frac{x_n}{y_n} = 1$, or, by Lemma 1, that $\lim_{n \rightarrow \infty} \frac{\Delta x_n}{\Delta y_n} = 1$.

We have

$$(37) \quad \begin{aligned} \lim_{n \rightarrow \infty} \frac{\Delta x_n}{\Delta y_n} &= \lim_{n \rightarrow \infty} (n a_n)^{-1} \left(\sum_{i=3}^n \frac{i a_i}{\lambda(i)} (\lambda(n) - \lambda(n-1)) + \frac{\lambda(n-1)}{\lambda(n)} n a_n \right) \\ &= 1 + \lim_{n \rightarrow \infty} \left(\frac{\lambda(n) - \lambda(n-1)}{n a_n} \sum_{i=3}^n \frac{i a_i}{\lambda(i)} \right). \end{aligned}$$

Note that, by (34), (36), we have

$$\sum_{i=3}^n \frac{i a_i}{\lambda(i)} \leq C \frac{n}{\lambda(n)} \sum_{i=3}^n a_i = O\left(\frac{n^2 (\ln n)^{1-\varepsilon}}{\lambda(n)}\right),$$

or, using condition $1/a_n = O(1)$, we find

$$(38) \quad \frac{1}{n a_n} \sum_{i=3}^n \frac{i a_i}{\lambda(i)} = O\left(\frac{n (\ln n)^{1-\varepsilon}}{\lambda(n)}\right).$$

Furthermore,

$$\lambda(n) - \lambda(n-1) \sim 2 \ln n / n \ln \ln n = 2 \frac{\lambda(n)}{n \ln n}.$$

Consequently, by (38), we have

$$\frac{\lambda(n) - \lambda(n-1)}{na_n} \sum_{i=3}^n \frac{ia_i}{\lambda(i)} = O\left(\frac{1}{\ln^\varepsilon n}\right)$$

and the lemma follows from (37). ■

Putting now, for $x = 2n$, $n \geq 3$,

$$x_n = \sum_{i=3}^n c(2i), \quad y_n = \sum_{i=3}^n 2i \ln \ln 2i / \ln^2 2i \prod_{p|i, p \text{ odd prime}} \frac{p-1}{p-2}.$$

Then, by the condition, $\lim_{n \rightarrow \infty} \frac{x_n - x_{n-1}}{y_n - y_{n-1}}$ exists and, by Lemma 1, it is sufficient to prove that

$$(39) \quad \lim_{n \rightarrow \infty} \frac{x_n}{y_n} = 2C_{tw}.$$

Putting

$$a_n = \prod_{p|i, p \text{ odd prime}} \frac{p-1}{p-2},$$

note that

$$a_n = \prod_{p|i, p \text{ odd prime}} \left(1 + \frac{1}{p-2}\right) = O(\ln \ln n)$$

and the conditions of Lemma 2 are satisfied. Using Theorem 5 and Lemma 2, we have

$$(40) \quad \lim_{n \rightarrow \infty} \frac{x_n}{y_n} = \lim_{n \rightarrow \infty} \frac{n^2}{\sum_{i=3}^n i \prod_{p|i, p \text{ odd prime}} \left(1 + \frac{1}{p-2}\right)}.$$

Furthermore,

$$(41) \quad \sum_{i=3}^n i \prod_{p|i, p \text{ odd prime}} \left(1 + \frac{1}{p-2}\right) = \sum_{i=3}^n i \sum_{d|i} \frac{\eta(d)}{d^*},$$

where $\eta(n) = 1$, if n is an odd square-free number, and $\eta(n) = 0$, otherwise;

$$d^* = \prod_{p|d} (p-2)^{\alpha_p}, \quad \text{if } d = \prod_{p|d} p^{\alpha_p}.$$

Continuing transformations of expression (41), we have

$$\begin{aligned} \sum_{i=3}^n i \prod_{p|i, p \text{ odd prime}} \left(1 + \frac{1}{p-2}\right) &= \sum_{d \leq n} \frac{\eta(d)}{d^*} \sum_{l \leq \frac{n}{d}} ld \\ &= \sum_{d \leq n} \frac{\eta(d)d}{d^*} \sum_{l \leq \frac{n}{d}} l = \sum_{d \leq n} \frac{\eta(d)d}{d^*} \frac{n^2}{2d^2} + O\left(n \sum_{d \leq n} \frac{1}{d}\right) \end{aligned}$$

$$\begin{aligned}
&= \frac{n^2}{2} \sum_{i=3}^n \frac{\eta(i)}{i^* i} + O(n \ln n) \\
&= \frac{n^2}{2} \prod_{p \geq 3, p \in P} \left(1 + \frac{1}{p(p-2)}\right) + O(n \ln n) \\
(42) \quad &= \frac{n^2}{2} \prod_{p \geq 3, p \in P} \left(1 - \frac{1}{(p-1)^2}\right)^{-1} + O(n \ln n) = \frac{n^2}{2C_{tw}} + O(n \ln n),
\end{aligned}$$

and (39) follows from (42) and (40). ■

7. CONCLUSIVE REMARKS

In this paper we selected only most interesting, in our opinion, binary partitions of positive integers. General results, which were obtained in [17], allow, by a similar way, to study many another binary partitions as well. Only for example, it could be proved that if, for $x \equiv 1 \pmod{4}$, $q(x) \geq 0$ denote the number of unordered representations of x by a sum of two squares of nonnegative integers, then

$$(43) \quad \sum_{i=0}^{\frac{x-1}{4}} q(4i+1) = \frac{\pi}{16}x + O(\sqrt{x}).$$

This formula follows from the exact equality (sf [17, example 4])

$$\begin{aligned}
\sum_{i=0}^{\frac{x-1}{4}} q(4i+1) &= \sum_{k=1}^{\lfloor (1+\sqrt{x})/2 \rfloor} \lfloor (1 + \sqrt{2x - (2k-1)^2})/2 \rfloor \\
(44) \quad &\quad - \left(\lfloor (1 + \sqrt{x})/2 \rfloor \right).
\end{aligned}$$

In addition, note that the last formula could be rewrite in the form

$$(45) \quad \sum_{i=0}^n q(4i+1) = \sum_{1 \leq k \leq r(n)} r(2n - k^2 + k) - \binom{r(n)}{2},$$

where $r(n)$ is the nearest integer to $\sqrt{n+1}$ and $n = \frac{x-1}{4}$.

REFERENCES

- [1] N. E. Bach, J. Shallit, *Algorithmic Number Theory*, MIT Press, 233(1996). ISBN 0-262-02405-5.
- [2] J. R. Chen, *On the representation of a large even integer as the sum of a prime and the product of at most two primes*, Kexue Tongbao, 17(1966), 385–386.
- [3] J. R. Chen, *On the representation of a larger even integer as the sum of a prime and the product of at most two primes*, Sci. Sinica, 16(1973), 157–176.
- [4] Y. Cai, *A remark on Chen's theorem*, Acta Arithm., 102, no.4 (2002), 339–352.
- [5] P. Dusart, *Autour de la fonction qui compte le nombre de nombres premiers*, Doctoral thesis for l'Universite de Limoges (1998).

- [6] P. Dusart , *The k -th prime is greater than $k(\ln k + \ln \ln k - 1)$ for $k \geq 2$* , Math. Comp. 68(1999), 411–415.
- [7] G. M. Fichtengolz, *Course of differential and integral calculus*, vol. I, Moscow, Nauka, 1969 (Russian).
- [8] G. Giordano, *On the irregularity of the distribution of the sums of pairs of odd primes*, International Journal of Mathematics and Mathematical Sciences, 30:6 (2002), 377–381.
- [9] A. Granville, J. van de Lune, and H. J. J. te Ree, *Checking the Goldbach conjecture on a vector computer*, Number Theory and Application (R. A. Mollin, ed.), Kluwer Acad., 1989, 423–433.
- [10] G. H. Hardy , J. E. Littlewood, *Tauberian theorems concerning power series and Dirichlet's series whose coefficients are positive*, Proc. London Math., 13, ser.2 (1913/14), 174–192.
- [11] G. H. Hardy and J. E. Littlewood, *Some problems of 'partitio numerorum'. III: on the expression of a number as a sum of primes*, Acta Math. 44, 1–70, 1922.
- [12] P. Haukkanen, *On generalized Landau identities*, Portugal. Math. 52, 1 (1995), 29–38.
- [13] E. Landau, *Über die zahlentheoretische Funktion $\varphi(n)$ und ihre Beziehung zum Goldbachschen Satz*, Göttinger Nachrichten (1900), 177–186.
- [14] J. E. Littlewood, *Sur la distribution des nombres premiers*, C. R., 158 (1914), 1869–1872.
- [15] P. M. Ross, *On Chen's theorem that each large even number has the form $(p_1 + p_2)$ or $(p_1 + p_2 p_3)$* , J. London Math. Soc. 2 (1975), 500–506.
- [16] B. Rosser, *Explicit Bounds for some Functions of Prime Numbers*, Amer. J. Math. 63 (1941), 211–232.
- [17] V. Shevelev, *Binary additive problems: recursions for numbers of representations*, <http://www.arxiv.org/abs/0901.3102>.
- [18] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences* (<http://www.research.att.com>)
- [19] E. W. Weisstein, *"Semiprime"* From MathWorld—A Wolfram Web Resource (<http://mathworld.wolfram.com>)

DEPARTMENTS OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA 84105, ISRAEL. E-MAIL:SHEVELEV@BGU.AC.IL